SPECIMEN LABEL

Growforce™ Max Fruit

3-16-22

Guaranteed Analysis

o dianamino a minamino a		
Total Nitrogen	3.0%	
3.0% Urea Nitrogen		
Available Phosphate (P ₂ O ₅)	16.0%	
Soluble Potash (K ₂ O)	22.0%	
Sulfur (S)	1.3%	
Magnesium (Mg)	0.01%	

Derived from: Urea, Dipotassium phosphate, and Potassium thiosulfate, and Magnesium sulfate

KEEP OUT OF REACH OF CHILDREN



WARNING

Causes skin irritation. Causes serious eye irritation.

PRECAUTIONARY STATEMENTS

Wash hands thoroughly after handling. Wear protective gloves and eye protection.

FIRST AID

IF ON SKIN: Wash with plenty of soap and water. If skin irritation occurs: Get medical advice/attention. Take off contaminated clothing and wash it before reuse.

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/attention.

STORAGE AND DISPOSAL

Dispose of contents/container in accordance with Federal, state and local laws and regulations.

COMPATIBILITY

Growforce Max Fruit is compatible with a wide range of commonly used agricultural chemicals. When mixing with other chemicals, always mix a small quantity and check for precipitate (fall out) in the mixture. If precipitation occurs, DO NOT proceed with application. If the mixture does not precipitate and it's the first time you have used this combination of products, test a small area of the crop prior to treatment of the whole area to test for phytotoxicity.

MIXING INSTRUCTIONS

Always fill the spray tank to 50% capacity with water before adding any tank mix agent, chemicals or other fertilizers. Add the full required amount of **Growforce Max Fruit** to the apparatus and add remaining water. Agitation is recommended during the mixing process.

GENERAL INFORMATION

Growforce Max Fruit is designed to provide a high level of potassium to aid in flowering and fruit fill. For best results apply **Growforce Max Fruit** during general irrigation.

DIRECTIONS FOR USE

Foliar and/or Fertigation Application:

rollal allu/ol	Foliar and/or Fertigation Application:		
Crop	Rate/ Acre	Comments	
Avocados, Mangos and Other Tropical Fruit	4 – 9 qts (Fertigation)	Apply at 7 - 14 day intervals from fruit set through to harvest.	
Berry Fruits	1 – 2 qts (Foliar) 1.5 – 3 qts (Fertigation)	Apply every 7 - 10 days from flowering through to harvest.	
Brassicas	2 – 4 qts (Foliar) 4 – 6 qts (Fertigation)	Apply every 14 days from head formation to harvest.	
Broadacre Crops (Not otherwise listed)	2 – 4 qts	Apply at regular intervals at vegetative and flowering to fruiting.	
Carrots	1.5 – 3 qts (Foliar) 2 – 4 qts (Fertigation)	Apply every 7 - 14 days once sufficient leaf area.	
Cereals	1.5 – 3 qts	Apply post emergence well before flowering.	
Pastures	1.5 – 3 qts	Apply post nodulation at regular intervals or to boost P & K.	
Cucurbits	2 – 4 qts (Foliar) 2 – 6 qts (Fertigation)	Apply every 7 - 14 days from flowering to harvest.	
Citrus	3 – 4 qts (Fertigation)	Apply from fruit set through to harvest every 14 days.	
Leafy greens	1.5 – 3 qts (Foliar) 3 – 5 qts (Fertigation)	Every 7 - 14 days from mid-crop through to harvest.	
Legumes	1 – 2 qts (Foliar) 3 – 6 qts (Fertigation)	Apply at flowering and during pod development.	
Macadamias	2 – 3 qts (Foliar) 4 – 8 (Fertigation)	Apply from nut development through to harvest at 14 - 21 day intervals.	
Olives	3 – 4 qts (Foliar) 4 – 6 qts (Fertigation)	Apply at 7 - 14 day intervals from fruit set through to harvest.	
Onions	1.5 – 2 qts (Foliar) 2 – 4 qts (Fertigation)	Apply from bulb formation to harvest.	
Pineapples	2 – 4 qts (Fertigation)	Apply at 14 - 21 day intervals from fruit set through to harvest.	
Pome Fruit	4 – 6 qts (Fertigation)	Apply every 7 - 14 days from fruit set.	
Potatoes	1.5 – 2 qts (Foliar) 2 – 4 qts (Fertigation)	Apply every 7 - 14 day from tuber initiation to harvest.	
Stone Fruit	4 – 8 qts (Fertigation)	Apply at 7 - 14 day intervals from fruit set through to harvest.	
Strawberries	1.5 – 2 qts (Foliar) 4 – 6 qts (Fertigation)	Apply every 7 - 10 days from flowering to harvest.	
Tomatoes	2 – 3 qts (Foliar) 4 – 6 qts (Fertigation)	Apply every 7 - 10 days from second truss formation to harvest.	
Vines: Table & Wine Grapes	1 – 2 qts (Foliar) 4 – 6 qts (Fertigation)	Apply every 14 days from flowering onwards. Do not exceed 2x concentration or 2x per acre rate.	

NET CONTENTS: 2.5 GAL (9.46 L)
WEIGHT PER GALLON: 11.85 lbs. (5.38 Kg)
NET WEIGHT: 29.63 lbs. (13.44 Kg)

SPECIMEN LABEL

WARRANTY DISCLAIMER

Vivid Life Sciences, LLC warrants only that the chemical composition of this product conforms to the ingredient statement given on the label and that the product is reasonably suited for the labeled use when applied according to the Directions for Use. Vivid Life Sciences, LLC makes no other warranties or representations of any kind, express or implied, concerning the product, including no implied warranty of merchantability or fitness for any particular purpose, and no such warranty shall be implied by law.

REV051617

NET CONTENTS: 2.5 GAL (9.46 L) WEIGHT PER GALLON: 11.85 lbs. (5.38 Kg) NET WEIGHT: 29.63 lbs. (13.44 Kg)