# **Growforce<sup>®</sup> Agri Max**

# 13-8-11

### **Guaranteed Analysis**

Total Nitrogen (N)	13.0%
3.0% Ammoniacal Nitrogen	
10.0% Urea Nitrogen	
Available Phophate (P <sub>2</sub> O <sub>5</sub> )	8.0%
Soluble Potash (K <sub>2</sub> O)	11.0%
Sulfur (S)	1.5%
1.5% Combined Sulfur (S)	
Molybdenum (Mo)	0.001%

Derived from: Urea, Monopotassium phosphate, Potassium thiosulfate, Potassium humate, Diammonium phosphate, Sodium molybdate dihydrate

Shake well before using.

# **KEEP OUT OF REACH OF CHILDREN**



Causes serious eye irritation. Causes skin irritation.

#### PRECAUTIONARY STATEMENTS

Wash hands thoroughly after handling. Wear eye protection and protective gloves.

#### FIRST AID

**IF IN EYES:** Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/attention.

**IF ON SKIN:** Wash with plenty of water and soap. No specific treatment available. If skin irritation occurs: Get medical advice/attention. Take off contaminated clothing and wash it before reuse.

#### STORAGE AND DISPOSAL

Dispose of contents/container in accordance with Federal, state and local laws and regulations.

#### COMPATIBILITY

**Growforce Agri-Max** is compatible with a wide range of commonly used agricultural chemicals. When mixing with other chemicals, always mix a small quantity and check for precipitate (fall out) in the mixture. If precipitation occurs, DO NOT proceed with application. If the mixture does not precipitate and it's the first time you have used this combination of products, test a small area of the crop prior to treatment of the whole crop.

#### WARRANTY DISCLAIMER

Vivid Life Sciences, LLC warrants only that the chemical composition of this product conforms to the ingredient statement given on the label and that the product is reasonably suited for the labeled use when applied according to the Directions for Use. Vivid Life Sciences, LLC makes no other warranties or representations of any kind, express or implied, concerning the product, including no implied warranty of merchantability or fitness for any particular purpose, and no such warranty shall be implied by law.

# **DIRECTIONS FOR USE**

# Foliar and Fertigation Application:

Crop	Rate/ Acre	Comments
Avocados, Mangoes and Other Tropical Fruit	1 – 2 gallons	Fertigation: Apply at 7 - 14 day intervals through to fruit set only
Cereals	3 – 4 pints	Foliar: Apply post emergence well before flowering
Broadacre Crops – not otherwise listed	4 – 6 pints	Foliar: Apply at regular intervals at vegetative , flowering and seed set stages
Cotton	4 – 5 pints	For cotton, apply to boost growth early in the season and post waterlogging events if required to boost growth
Legumes	4 – 5 pints	In legumes, apply at regular intervals from 20 days after establishment until onset of nodulation
Pastures	3 – 6 pints	Apply at regular intervals to boost growth
Citrus	4 – 6 pints	Foliar: Spray as and when required as per tissue tests
	1 – 2 gallons	Fertigation: Spray as and when required as per tissue tests
Cucurbits	4 – 6 pints	Foliar: Spray as and when required as per tissue tests
	1 – 2 gallons	Fertigation: Spray as and when required as per tissue tests
Leafy Greens	3 – 7 pints	Foliar: Apply every 7 - 14 days from mid-crop through to harvest
	7 – 10 pints	Fertigation: Apply every 7 - 14 days from mid-crop through to harvest
Olives	6 – 8.5 pints	Foliar: Apply at 7 - 14 day intervals before fruit set
	8.5 – 13 pints	Fertigation: Apply at 7 - 14 day intervals before fruit set
Onions, Potatoes	6 – 8.5 pints	Foliar: Apply from bulb formation to harvest in onions and from hooking to canopy closure in potatoes
	8.5 – 13 pints	Fertigation: Apply from bulb formation to harvest in onions and from hooking to canopy closure in potatoes
Stone & Pome Fruit	1 – 2 gallons	Fertigation: Apply at 7 - 14 day intervals from fruit set through to harvest
Tomatoes	4 – 5 pints	Foliar: Apply every 7 - 10 days from planting to 4th truss formation
	1 – 2.7 pints	Fertigation: Apply every 7 - 10 days from planting to 4th truss formation
Vines	4 pints or 10 – 34 fl. oz./ 34 fl. oz. of water	Foliar: Apply every 14 days from flowering onwards. Do not exceed 2x concentration or 2x per hectare rate.
Table or Wine grapes	1 – 2 gallons	Fertigation: Apply the higher fertigated rate post-harvest
	1 – 2 gallons	<b>o</b> 11 <i>9</i> <b>o</b> 0

REV102017

#### NET CONTENTS: 2.5 GAL (9.46 L) WEIGHT PER GALLON: 11.18 lbs. (5.07 Kg) NET WEIGHT: 27.95 lbs. (12.68 Kg)

F002511 Manufactured for and Guaranteed By: Vivid Life Sciences, LLC, P.O. Box 41632, Minneapolis, MN 55441